



Albac[®] 10%, 15% Poult

Proven solution for Necrotic Enteritis
authenticated by Huvepharma

PRODUCT DESCRIPTION

Albac® 10%, contains feed grade zinc bacitracin equivalent to 100 grams (10%) bacitracin per kilogram.
 Albac® 15%, contains feed grade zinc bacitracin equivalent to 150 grams (15%) bacitracin per kilogram.

INDICATIONS FOR USE

1. For the prevention and control of necrotic enteritis in chickens caused or complicated by *Clostridium spp*
2. Increased weight gain and improved feed efficiency.
3. Increased egg production and improved feed efficiency in laying hens.

KEY POINTS

- Albac® helps lowering intestinal counts of *C. perfringens* as well as other pathogenic bacteria, this promotes a healthy gut flora and higher nutrient absorption in broilers.
- The only bacitracin available for use in all ages of laying hens.
- Not absorbed from the intestinal tract, thus no withdrawal period is required prior to slaughter.
- Consistent performance after long time use.
- Effective in extreme weather conditions.


USERS INFORMATION: ALBAC®

Active Drug Ingredient		Each kilogram contains feed grade zinc bacitracin.		
For Use in Manufactured Medicated Feeds				
Species	Grams bacitracin per tonne of Feed	Albac®10% dosage	Albac®15% dosage	Indications for Use
Broiler chickens	25	dose of 250 gram /mtion	dose of 170 gram /mtion	For increased weight gain and improved feed efficiency
	50	dose of 500 gram/mtion	dose of 333 gram/mtion	Supporting the prevention of necrotic enteritis. Medicated feed should be fed to poultry during the period when they are at most risk of the disease, which is usually between 2 and 12 weeks.
	200	dose of 2000 gram/mtion	dose of 1333 gram/mtion	Supporting the treatment of necrotic enteritis: add Albac® for 5-7 days.
Laying hens	50-75	dose of 500-750 gram/mtion	dose of 333-500 gram/mtion	For increased weight gain and improved feed efficiency

Can be used in combination with our registered feed additives and coccidiostats*.

HANDLING

- Albac® is packaged in a 25 kg, multilayer paper bag with protective barrier ply.
- Store in a cool (<25°C), dry place and keep package closed to avoid contamination.
- Shelf life is 36 months from the date of manufacturing.
- For use in dry feeds only.
- Zero day withdrawal period.



*Specific claims vary by country.

Please contact your Huvepharma representative or distributor for more information.

References available on request

Huvepharma N.V.
 Uitbreidingsstraat 80 • 2600 Antwerp • Belgium
 P +32 3 288 1849 • F +32 3 289 7845
 customerservice@huvepharma.com


HUVEPHARMA®
 We add performance to your business
<http://www.huvepharma.com>

© 2015 Huvepharma EAD. All right reserved

Huvepharma SEA (Pune) Pvt. Ltd.
 104, 1st Floor, 'Green Terraces', Lane No.5
 South Koregaon Park, Kawdewadi, Pune 411028
 P +91 20 2615 4193 • F +91 20 2615 3973
 salesindia@huvepharma.com