

THE NEW COMBINATION COCCIDIOSTAT FOR BROILERS




PHARMACOLOGICAL ACTION

Nimax® is a microGranulated coccidiostat containing 2 active ingredients. The combined strength of the chemical **nicarbazin** and the ionophore **maduramicin** leads to a unique new product in coccidiosis control.

Nimax® is effective against coccidiosis caused by all major *Eimeria* species in broilers. On cellular level the effect of Nimax® to inhibit the development of sporozoites and trophozoites is enhanced by the **synergistic** action of nicarbazin and maduramicin.

This innovative combination product will not only result in **superior coccidiosis control**, but will also **enhance performance levels**.



● nicarbazin: 160 g/kg
● maduramicin: 15 g/kg

 **HUVEPHARMA®**
We add performance to your business



COMPOSITION

| Content | NIMAX® |
|-------------|-----------------|
| Nicarbazin | 16 g per 100 g |
| Maduramicin | 1.5 g per 100 g |

INDICATIONS

As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. acervulina*, *E. maxima*, *E. necatrix*, *E. brunetti*, *E. mitis* and *E. praecox* in chickens.

CONTRAINDICATIONS

Do not feed to horses or other equines. Do not mix with other coccidiostats.

TARGET SPECIES

Broiler chickens.

MODE OF ADMINISTRATION

Orally, well homogenized into the feed. Nimax® should be thoroughly and evenly mixed in the feed in accordance with current good manufacturing practice for medicated feed.

DOSAGES

For prevention of coccidiosis in broilers.

Nimax® is used at an inclusion rate of **250 g per ton of feed**

| | | Broiler chickens |
|--|---------|------------------|
| Content of nicarbazin/maduramicin in complete feedingstuff (ppm) | maximum | 40/3.75 |
| Quantity of Nimax® incorporated in feed (g per ton) | maximum | 250 |

INSTRUCTIONS FOR USE

Nimax® microgranules must be mixed with other ingredients before feeding in order to have an optimal homogeneity and to avoid segregation. Mixing and conveying equipment should be properly cleaned to prevent carry-over.

SIDE EFFECTS

None under correct use of the product.

WITHDRAWAL PERIOD

None under correct use of the product.

STORAGE

In the original packing, well closed, in dry and well-ventilated facilities, protected from direct sunlight.

PACKING

Nimax® is packed in 25 kg bags.

WARNING

Dangerous for equines.

When mixing, direct contact with the skin and mucous membranes should be avoided. Accidental ingestion should be avoided. Wear overalls, safety glasses, mask and impervious gloves when handling or mixing the product. Contaminated clothing should be removed and any splashes on to the skin should be washed off immediately. If accidental eye contact occurs, immediately rinse thoroughly with water. Seek medical advice if irritation persists. Wash hands after use.

Huvepharma N.V.
Uitbreidingsstraat 80 • 2600 Antwerp • Belgium
P +32 3 288 1849 • F +32 3 289 7845
customerservice@huvepharma.com


<http://www.huvepharma.com>

Huvepharma SEA (Pune) Pvt. Ltd.
104, 1st Floor, 'Green Terraces', Lane No.5
South Koregaon Park, Kawdewadi, Pune 411028
P +91 20 2615 4193 • F +91 20 2615 3973
salesindia@huvepharma.com

© 2015 Huvepharma EAD. All right reserved